

EASTERN POLLINATOR MIXTURE

Wildflower Mixture*

Common Name	Scientific Name
Lavender Hyssop	Agastache foeniculum
Butterfly Milkweed	Asclepias tuberosa
New England Aster	Aster novae-angliae
Blue Wild Indigo	Baptisia australis
False Boneset	Brickellia eupatorioides
Partridge Pea	Chamaecrista fasciculata
Lance-Leaved Coreopsis	Coreopsis lanceolata
Plains Coreopsis	Coreopsis tinctoria
Showy Tick Trefoil	Desmodium canadense
Purple Coneflower	Echinacea purpurea
Indian Blanket	Gaillardia pulchella
Perennial Lupine	Lupinus perennis
Lemon Mint	Monarda citriodora
Spotted Beebalm	Monarda punctata
Evening Primrose	Oenothera biennis
Hairy Penstemon	Penstemon hirsutus
Slender Mountain Mint	Pycnanthemum tenuifolium
Black Eyed Susan	Rudbeckia hirta
Brown-Eyed Susan	Rudbeckia triloba
Gray Goldenrod	Solidago nemoralis
Ohio Spiderwort	Tradescantia ohioensis

Seeds/lb. = 316,000

Planting rate = 8-16 lbs./acre, 6 oz./1000 sq. feet

Planting times: Best times to plant are spring, early summer and fall.

Planting range: Formulated for the eastern United States and southeast Canada where pollinator conservation is desired. The western boundary would be a line straight south from the eastern borders of North and South Dakota.

*Mixture may vary occasionally from the indicated listing, based on availability of individual species.